

BRENTWOOD PHARMACY
761 Suffolk Ave , Brentwood , NY , 11717
Phone#: 6312733314 Fax#: 6312738863

Drug NDC: 00074-6799-22
Drug Name: KALETRA

How should I use this medicine?

Take this medicine by mouth with a glass of water. Follow the directions on the prescription label. You can take this medicine with or without food. Do not crush, break, or chew the tablets. Take your medicine at regular intervals. Do not take your medicine more often than directed. For your anti-HIV therapy to work as well as possible, take each dose exactly as prescribed. Do not skip doses or stop your medicine even if you feel better. Skipping doses may make the HIV virus resistant to this medicine and other medicines. Do not stop taking except on your doctor's advice. A special MedGuide will be given to you by the pharmacist with each prescription and refill. Be sure to read this information carefully each time. Talk to your pediatrician regarding the use of this medicine in children. While this drug may be prescribed for selected conditions, precautions do apply.

Overdosage: If you think you have taken too much of this medicine contact a poison control center or emergency room at once.
NOTE: This medicine is only for you. Do not share this medicine with others.

Where should I keep my medicine?

Keep out of the reach of children. Store at room temperature between 15 and 30 degrees C (59 and 86 degrees F). Keep in original container. Do not store this medicine outside the original container for longer than 2 weeks. Protect from moisture and humidity. Throw away any unused medicine after the expiration date.

NOTE: This sheet is a summary. It may not cover all possible information. If you have questions about this medicine, talk to your doctor, pharmacist, or health care provider.

What should I watch for while using this medicine?

Visit your doctor or health care professional for regular check ups. Discuss any new symptoms with your doctor. You will need to have important blood work done while on this medicine. HIV is spread to others through sexual or blood contact. Talk to your doctor about how to stop the spread of HIV. Birth control pills may not work properly while you are taking this medicine. Talk to your doctor about using an extra method of birth control. Women who can still have children must use a reliable form of barrier contraception, like a condom or diaphragm. This medicine may affect blood sugar levels. If you have diabetes, check with your doctor or health care professional before you change your diet or the dose of your diabetic medicine.

What is this medicine?

LOPINAVIR; RITONAVIR (loe PIN a veer; ri TOE na veer) is two antiretroviral medicines in one tablet. It is used with other medicines to treat HIV. This medicine is not a cure for HIV. It will not stop the spread of HIV to others.

This medicine may be used for other purposes; ask your health care provider or pharmacist if you have questions.

What should I tell my health care provider before I take this medicine?

They need to know if you have any of these conditions: diabetes -hemophilia -liver disease -an unusual or allergic reaction to lopinavir, ritonavir, other medicines, foods, dyes, or preservatives -pregnant or trying to get pregnant -breast-feeding

What if I miss a dose?

If you miss a dose, take it as soon as you can. If it is almost time for your next dose, take only that dose. Do not take double or extra doses.

What side effects may I notice from receiving this medicine?

Side effects that you should report to your doctor or health care professional as soon as possible: allergic reactions like skin rash, itching or hives, swelling of the face, lips, or tongue -fast, irregular heart rate -feeling faint or lightheaded, falls -nausea, vomiting, abdominal pain -red, hot, painful spot on leg -redness, blistering, peeling or loosening of the skin, including inside the mouth Side effects that usually do not require medical attention (report to your doctor or health care professional if they continue or are bothersome): diarrhea -difficulty sleeping -headache -heartburn -increased hunger or thirst -increased urination -unusually weak or tired -weight gain around waist, back, or thinning of face, arms, legs

This list may not describe all possible side effects. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.