

BRENTWOOD PHARMACY
761 Suffolk Ave , Brentwood , NY , 11717
Phone#: 6312733314 Fax#: 6312738863

Drug NDC: 00093-1015-01
Drug Name: NABUMETONE

How should I use this medicine?

Take this medicine by mouth with a full glass of water. Follow the directions on the prescription label. You can take it with or without food. If it upsets your stomach, take it with food. Try to not lie down for at least 10 minutes after you take this medicine. Take your medicine at regular intervals. Do not take your medicine more often than directed. Long term, continuous use may increase the risk of heart attack or stroke. A special MedGuide will be given to you by the pharmacist with each prescription and refill. Be sure to read this information carefully each time. Talk to your pediatrician regarding the use of this medicine in children. Special care may be needed.

Overdosage: If you think you have taken too much of this medicine contact a poison control center or emergency room at once. NOTE: This medicine is only for you. Do not share this medicine with others.

Where should I keep my medicine?

Keep out of the reach of children. Store at room temperature between 15 and 30 degrees C (59 and 86 degrees F). Keep container tightly closed. Throw away any unused medicine after the expiration date.

NOTE: This sheet is a summary. It may not cover all possible information. If you have questions about this medicine, talk to your doctor, pharmacist, or health care provider.

What should I watch for while using this medicine?

Tell your doctor or health care professional if your pain does not get better. Talk to your doctor before taking another medicine for pain. Do not treat yourself. This medicine does not prevent heart attack or stroke. In fact, this medicine may increase the chance of a heart attack or stroke. The chance may increase with longer use of this medicine and in people who have heart disease. If you take aspirin to prevent heart attack or stroke, talk with your doctor or health care professional. Do not take medicines such as ibuprofen and naproxen with this medicine. Side effects such as stomach upset, nausea, or ulcers may be more likely to occur. Many medicines available without a prescription should not be taken with this medicine. This medicine can cause ulcers and bleeding in the stomach and intestines at any time during treatment. Do not smoke cigarettes or drink alcohol. These increase irritation to your stomach and can make it more susceptible to damage from this medicine. Ulcers and bleeding can happen without warning symptoms and can cause death. You may get drowsy or dizzy. Do not drive, use machinery, or do anything that needs mental alertness until you know how this medicine affects you. Do not stand or sit up quickly, especially if you are an older patient. This reduces the risk of dizzy or fainting spells. This medicine can cause you to bleed more easily. Try to avoid damage to your teeth and gums when you brush or floss your teeth.

What is this medicine?

NABUMETONE (na BYOO me tone) is a non-steroidal anti-inflammatory drug (NSAID). It is used to reduce swelling and to treat pain. It is used for osteoarthritis or rheumatoid arthritis.

This medicine may be used for other purposes; ask your health care provider or pharmacist if you have questions.

What should I tell my health care provider before I take this medicine?

They need to know if you have any of these conditions:
-cigarette smoker
-coronary artery bypass graft (CABG) surgery within the past 2 weeks
-drink more than 3 alcohol-containing drinks a day
-heart disease
-high blood pressure
-history of stomach bleeding
-kidney disease
-liver disease
-lung or breathing disease, like asthma
-an unusual or allergic reaction to nabumetone, aspirin, other NSAIDs, other medicines, foods, dyes, or preservatives
-pregnant or trying to get pregnant
-breast-feeding

What if I miss a dose?

If you miss a dose, take it as soon as you can. If it is almost time for your next dose, take only that dose. Do not take double or extra doses.

What side effects may I notice from receiving this medicine?

Side effects that you should report to your doctor or health care professional as soon as possible:
-black or bloody stools, blood in the urine or vomit
-blurred vision
-chest pain
-difficulty breathing or wheezing
-nausea or vomiting
-skin rash, skin redness, blistering or peeling skin, hives, or itching
-slurred speech or weakness on one side of the body
-severe stomach pain
-swelling of eyelids, throat, lips
-unexplained weight gain or swelling
-unusually weak or tired
-yellowing of eyes or skin
Side effects that usually do not require medical attention (report to your doctor or health care professional if they continue or are bothersome):
-constipation or diarrhea
-gas or heartburn

This list may not describe all possible side effects. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.