

BRENTWOOD PHARMACY
761 Suffolk Ave , Brentwood , NY , 11717
Phone#: 6312733314 Fax#: 6312738863

Drug NDC: 00172-4333-70
Drug Name: NAFAZODONE 150MG

How should I use this medicine?

Take this medicine by mouth with a glass of water. Follow the directions on the prescription label. You may take this medicine with or without food. Take your medicine at regular intervals. Do not take your medicine more often than directed. Do not stop taking this medicine suddenly except upon the advice of your doctor. Stopping this medicine too quickly may cause serious side effects or your condition may worsen. A special MedGuide will be given to you by the pharmacist with each prescription and refill. Be sure to read this information carefully each time. Talk to your pediatrician regarding the use of this medicine in children. Special care may be needed.

Overdosage: If you think you have taken too much of this medicine contact a poison control center or emergency room at once. NOTE: This medicine is only for you. Do not share this medicine with others.

Where should I keep my medicine?

Keep out of the reach of children. Store at room temperature below 40 degrees C (104 degrees F). Keep container tightly closed. Throw away any unused medicine after the expiration date.

NOTE: This sheet is a summary. It may not cover all possible information. If you have questions about this medicine, talk to your doctor, pharmacist, or health care provider.

What should I watch for while using this medicine?

Tell your doctor if your symptoms do not get better or if they get worse. Visit your doctor or health care professional for regular checks on your progress. Because it may take several weeks to see the full effects of this medicine, it is important to continue your treatment as prescribed by your doctor. Patients and their families should watch out for new or worsening thoughts of suicide or depression. Also watch out for sudden changes in feelings such as feeling anxious, agitated, panicky, irritable, hostile, aggressive, impulsive, severely restless, overly excited and hyperactive, or not being able to sleep. If this happens, especially at the beginning of treatment or after a change in dose, call your health care professional. You may get drowsy or dizzy. Do not drive, use machinery, or do anything that needs mental alertness until you know how this medicine affects you. Do not stand or sit up quickly, especially if you are an older patient. This reduces the risk of dizzy or fainting spells. Alcohol may interfere with the effect of this medicine. Avoid alcoholic drinks. Your mouth may get dry. Chewing sugarless gum or sucking hard candy, and drinking plenty of water will help. Contact your doctor if the problem does not go away or is severe.

What is this medicine?

NEFAZODONE (nef AY zoe done) is used to treat depression.

This medicine may be used for other purposes; ask your health care provider or pharmacist if you have questions.

What should I tell my health care provider before I take this medicine?

They need to know if you have any of these conditions: bipolar disorder or mania; glaucoma; heart disease; liver disease; seizures; suicidal thoughts, plans, or attempt; a previous suicide attempt by you or a family member; an unusual or allergic reaction to nefazodone, other medicines, food, dyes, or preservatives; pregnant or trying to get pregnant; breast-feeding

What if I miss a dose?

If you miss a dose, take it as soon as you can. If it is almost time for your next dose, take only that dose. Do not take double or extra doses.

What side effects may I notice from receiving this medicine?

Side effects that you should report to your doctor or health care professional as soon as possible: allergic reactions like skin rash, itching or hives, swelling of the face, lips, or tongue; changes in vision; confusion; elevated mood, decreased need for sleep, racing thoughts, impulsive behavior; eye pain; feeling faint or lightheaded, falls; feeling agitated, angry, or irritable; fever or chills, sore throat; hallucination, loss of contact with reality; irregular heartbeat; loss of balance or coordination; painful or prolonged erections; redness, blistering, peeling or loosening of the skin, including inside the mouth; restlessness, pacing, inability to keep still; seizures; signs and symptoms of liver injury like dark yellow or brown urine; general ill feeling or flu-like symptoms; light-colored stools; loss of appetite; nausea; right upper belly pain; unusually weak or tired; yellowing of the eyes or skin; stiff muscles; suicidal thoughts or other mood changes; trouble passing urine or change in the amount of urine; trouble sleeping; vomiting. Side effects that usually do not require medical attention (report to your doctor or health care professional if they continue or are bothersome): change in sex drive or performance; constipation; dizziness; dry mouth; indigestion, stomach upset; nausea

This list may not describe all possible side effects. Call your doctor for medical advice about side effects.

You may report side effects to FDA at 1-800-FDA-1088.