

BRENTWOOD PHARMACY
761 Suffolk Ave , Brentwood , NY , 11717
Phone#: 6312733314 Fax#: 6312738863

Drug NDC: 00173-0880-25
Drug Name: TABLOID

How should I use this medicine?

Take this medicine by mouth with a glass of water. Follow the directions on the prescription label. Take your medicine at regular intervals. Do not take it more often than directed. Do not stop taking except on your doctor's advice. Talk to your pediatrician regarding the use of this medicine in children. While this drug may be prescribed for selected conditions, precautions do apply.

Overdosage: If you think you have taken too much of this medicine contact a poison control center or emergency room at once.
NOTE: This medicine is only for you. Do not share this medicine with others.

Where should I keep my medicine?

Keep out of the reach of children. Store at room temperature between 15 and 25 degrees C (59 and 77 degrees F). Protect from moisture. Keep container tightly closed. Throw away any unused medicine after the expiration date.

NOTE: This sheet is a summary. It may not cover all possible information. If you have questions about this medicine, talk to your doctor, pharmacist, or health care provider.

What should I watch for while using this medicine?

This drug may make you feel generally unwell. This is not uncommon, as chemotherapy can affect healthy cells as well as cancer cells. Report any side effects. Continue your course of treatment even though you feel ill unless your doctor tells you to stop. In some cases, you may be given additional medicines to help with side effects. Follow all directions for their use. Call your doctor or health care professional for advice if you get a fever, chills or sore throat, or other symptoms of a cold or flu. Do not treat yourself. This drug decreases your body's ability to fight infections. Try to avoid being around people who are sick. This medicine may increase your risk to bruise or bleed. Call your doctor or health care professional if you notice any unusual bleeding. Be careful brushing and flossing your teeth or using a toothpick because you may get an infection or bleed more easily. If you have any dental work done, tell your dentist you are receiving this medicine. Avoid taking products that contain aspirin, acetaminophen, ibuprofen, naproxen, or ketoprofen unless instructed by your doctor. These medicines may hide a fever. Do not become pregnant while taking this medicine. Women should inform their doctor if they wish to become pregnant or think they might be pregnant. There is a potential for serious side effects to an unborn child. Talk to your health care professional or pharmacist for more information. Do not breast-feed an infant while taking this medicine.

What is this medicine?

THIOGUANINE (6-TG) (thyeh GWAH neen) is a chemotherapy drug. It is used to treat some leukemias.

This medicine may be used for other purposes; ask your health care provider or pharmacist if you have questions.

What should I tell my health care provider before I take this medicine?

They need to know if you have any of these conditions: infection (especially virus infection such as chickenpox or herpes) liver disease low blood counts like low platelets, red blood cells, or white blood cells thiopurine methyltransferase (TPMT) deficiency an unusual or allergic reaction to thioguanine, other chemotherapy, other medicines, foods, dyes, or preservatives pregnant or trying to get pregnant breast-feeding

What if I miss a dose?

If you miss a dose, skip that dose unless your prescriber or health care professional tells you otherwise. Do not take double or extra doses. If you vomit after taking a dose, call your prescriber or health care professional for advice.

What side effects may I notice from receiving this medicine?

Side effects that you should report to your doctor or health care professional as soon as possible: allergic reactions like skin rash, itching or hives, swelling of the face, lips, or tongue low blood counts - this medicine may decrease the number of white blood cells, red blood cells and platelets. You may be at increased risk for infections and bleeding. signs of infection - fever or chills, cough, sore throat, pain or difficulty passing urine signs of decreased platelets or bleeding - bruising, pinpoint red spots on the skin, black, tarry stools, blood in the urine signs of decreased red blood cells - unusually weak or tired, fainting spells, lightheadedness breathing problems mouth sores right upper belly pain swelling of the ankles, feet, hands, or stomach yellowing of the eyes or skin water or weight gain Side effects that usually do not require medical attention (report to your doctor or health care professional if they continue or are bothersome): diarrhea loss of appetite nausea, vomiting

This list may not describe all possible side effects. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.