

**BRENTWOOD PHARMACY**  
**761 Suffolk Ave , Brentwood , NY , 11717**  
**Phone#: 6312733314 Fax#: 6312738863**

**Drug NDC:** 00378-6709-77  
**Drug Name:** RABEPRAZOLE SODIUM

**How should I use this medicine?**

Take this medicine by mouth. Swallow the tablets whole with a drink of water. Follow the directions on the prescription label. Do not crush, break, or chew. You may take this medicine with or without food. Take your medicine at regular intervals. Do not take more often than directed. A special MedGuide will be given to you by the pharmacist with each prescription and refill. Be sure to read this information carefully each time. Talk to your pediatrician regarding the use of this medicine in children. Special care may be needed.

**Overdosage: If you think you have taken too much of this medicine contact a poison control center or emergency room at once.**  
**NOTE: This medicine is only for you. Do not share this medicine with others.**

**Where should I keep my medicine?**

Keep out of the reach of children. Store at room temperature between 15 and 30 degrees C (59 and 86 degrees F). Protect from light and moisture. Throw away any unused medicine after the expiration date.

**NOTE: This sheet is a summary. It may not cover all possible information. If you have questions about this medicine, talk to your doctor, pharmacist, or health care provider.**

**What should I watch for while using this medicine?**

It can take several days before your stomach pain gets better. Check with your doctor or health care professional if your condition does not start to get better, or if it gets worse. You may need blood work done while you are taking this medicine. This medicine may cause a decrease in vitamin B12. You should make sure that you get enough vitamin B12 while you are taking this medicine. Discuss the foods you eat and the vitamins you take with your health care professional.

**What is this medicine?**

RABEPRAZOLE (ra BE pray zole) prevents the production of acid in the stomach. It is used to treat gastroesophageal reflux disease (GERD), certain ulcers, inflammation of the esophagus, and Zollinger-Ellison Syndrome.

**This medicine may be used for other purposes; ask your health care provider or pharmacist if you have questions.**

**What should I tell my health care provider before I take this medicine?**

They need to know if you have any of these conditions: liver disease, low levels of magnesium in the blood, lupus, an unusual or allergic reaction to rabeprazole, esomeprazole, omeprazole, lansoprazole, pantoprazole, other medicines, foods, dyes, or preservatives, pregnant or trying to get pregnant, breast-feeding

**What if I miss a dose?**

If you miss a dose, take it as soon as you can. If it is almost time for your next dose, take only that dose. Do not take double or extra doses.

**What side effects may I notice from receiving this medicine?**

Side effects that you should report to your doctor or health care professional as soon as possible: allergic reactions like skin rash, itching or hives, swelling of the face, lips, or tongue; bone, muscle or joint pain; breathing problems; chest pain or chest tightness; dark yellow or brown urine; dizziness; fast, irregular heartbeat; feeling faint or lightheaded; fever or sore throat; muscle spasm; palpitations; rash on cheeks or arms that gets worse in the sun; redness, blistering, peeling or loosening of the skin, including inside the mouth; seizures; stomach polyps; tremors; unusual bleeding or bruising; unusually weak or tired; yellowing of the eyes or skin. Side effects that usually do not require medical attention (report to your doctor or health care professional if they continue or are bothersome): constipation; diarrhea; dry mouth; headache; nausea

**This list may not describe all possible side effects. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.**