

**BRENTWOOD PHARMACY**  
**761 Suffolk Ave , Brentwood , NY , 11717**  
**Phone#: 6312733314 Fax#: 6312738863**

**Drug NDC: 00603-0851-94**  
**Drug Name: Q-TAPP**

**How should I use this medicine?**

Take this medicine by mouth. Follow the directions on the prescription label. Use a specially marked spoon or container to measure each dose. Ask your pharmacist if you do not have one. Household spoons are not accurate. You can take it with or without food. If it upsets your stomach, take it with food. Take your medicine at regular intervals. Do not take it more often than directed. Talk to your pediatrician regarding the use of this medicine in children. While this drug may be prescribed for selected conditions, precautions do apply. Patients over 65 years old may have a stronger reaction and need a smaller dose.

**Overdosage: If you think you have taken too much of this medicine contact a poison control center or emergency room at once. NOTE: This medicine is only for you. Do not share this medicine with others.**

**Where should I keep my medicine?**

Keep out of the reach of children. This medicine may cause accidental overdose and death if taken by other adults, children, or pets. Mix any unused medicine with a substance like cat litter or coffee grounds. Then throw the medicine away in a sealed container like a sealed bag or a coffee can with a lid. Do not use the medicine after the expiration date. Store between 8 and 30 degrees C (46 and 86 degrees F). Do not freeze. Protect from light.

**NOTE: This sheet is a summary. It may not cover all possible information. If you have questions about this medicine, talk to your doctor, pharmacist, or health care provider.**

**What should I watch for while using this medicine?**

Visit your doctor or health care professional for regular check ups. Tell your doctor if your symptoms do not improve or if they get worse. Your mouth may get dry. Chewing sugarless gum or sucking hard candy, and drinking plenty of water may help. Contact your doctor if the problem does not go away or is severe. This medicine may cause dry eyes and blurred vision. If you wear contact lenses you may feel some discomfort. Lubricating drops may help. See your eye doctor if the problem does not go away or is severe. You may get drowsy or dizzy. Do not drive, use machinery, or do anything that needs mental alertness until you know how this medicine affects you. Do not stand or sit up quickly, especially if you are an older patient. This reduces the risk of dizzy or fainting spells. Alcohol may interfere with the effect of this medicine. Avoid alcoholic drinks.

**What is this medicine?**

BROMPHENIRAMINE; PSEUDOEPHEDRINE (brome fen IR a meen; soo doe e FED rin) is a combination of an antihistamine and a decongestant. It is used to treat the symptoms of allergy and colds. This medicine will not treat an infection.

**This medicine may be used for other purposes; ask your health care provider or pharmacist if you have questions.**

**What should I tell my health care provider before I take this medicine?**

They need to know if you have any of the following conditions: asthma, diabetes, difficulty passing urine, glaucoma, heart disease, high blood pressure, taken an MAOI like Carbox, Eldepryl, Marplan, Nardil, or Parmate in last 14 days, thyroid disease, ulcer or other stomach problem, an unusual or allergic reaction to brompheniramine, pseudoephedrine, other medicines, foods, dyes, or preservatives, pregnant or trying to get pregnant, breast-feeding

**What if I miss a dose?**

If you miss a dose, take it as soon as you can. If it is almost time for your next dose, take only that dose. Do not take double or extra doses.

**What side effects may I notice from receiving this medicine?**

Side effects that you should report to your doctor or health care professional as soon as possible: allergic reactions like skin rash, itching or hives, swelling of the face, lips, or tongue; anxiety, fear; change in vision; cold, clammy skin; confused, faint, lightheaded; difficulty passing urine; fast or irregular heartbeat; hallucinations; high blood pressure; seizure; tremor; trouble breathing; vomiting; Side effects that usually do not require medical attention (report to your doctor or health care professional if they continue or are bothersome): headache; loss of appetite; nervous, restless; stomach upset, nausea; trouble sleeping

**This list may not describe all possible side effects. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.**