

BRENTWOOD PHARMACY
761 Suffolk Ave , Brentwood , NY , 11717
Phone#: 6312733314 Fax#: 6312738863

Drug NDC: 23594-0780-88
Drug Name: ULESFIA 2-PACK

How should I use this medicine?

This medicine is for external use only. Do not take by mouth. Follow the directions on the product label. Do not wash hair with conditioner or a combination shampoo-conditioner immediately before applying this medicine. Follow product-specific instructions for length of application and product removal. All products should be rinsed from the hair over a sink or tub to limit skin exposure; do not use hot water. In general, do not re-wash the hair for 1 to 2 days after use. Talk to your pediatrician regarding the use of this medicine in children. While this medicine may be prescribed for children as young as 6 months for selected conditions, precautions do apply.

Overdosage: If you think you have taken too much of this medicine contact a poison control center or emergency room at once. NOTE: This medicine is only for you. Do not share this medicine with others.

Where should I keep my medicine?

Keep out of the reach of children. Store at room temperature between 15 and 30 degrees C (59 and 86 degrees F). Do not freeze. Throw away any unused medicine after the expiration date.

NOTE: This sheet is a summary. It may not cover all possible information. If you have questions about this medicine, talk to your doctor, pharmacist, or health care provider.

What should I watch for while using this medicine?

All recently worn clothing, hats, used bedding, and towels should be washed in hot water or dry-cleaned. Combs, brushes, and hair clips should also be washed in hot water. This medicine gets rid of lice, but it does not get rid of their eggs. A second treatment is needed 7 days after the first treatment. Head lice can be spread from one person to another by direct contact with clothing, hats, scarves, bedding, towels, washcloths, hairbrushes, and combs. All members of your household should be examined for head lice and should receive treatment if they are found to be infected. If you have any questions about this, check with your doctor or health care professional.

What is this medicine?

BENZYL ALCOHOL (BEN zil AL kuh hawl) lotion is used to treat lice of the hair and scalp. It acts by destroying the lice. It does not destroy their eggs; a second treatment is needed 7 days after the first treatment.

This medicine may be used for other purposes; ask your health care provider or pharmacist if you have questions.

What should I tell my health care provider before I take this medicine?

They need to know if you have any of these conditions: an unusual or allergic reaction to benzyl alcohol, other medicines, foods, dyes, or preservatives; breast feeding; pregnant or trying to get pregnant

What if I miss a dose?

If you miss your second dose, use it as soon as you can. Do not use double or extra doses.

What side effects may I notice from receiving this medicine?

Side effects that you should report to your doctor or health care professional as soon as possible: allergic reactions like skin rash, itching or hives, swelling of the face, lips, or tongue; eye or skin irritation; Side effects that usually do not require medical attention (report to your doctor or health care professional if they continue or are bothersome): itching; redness

This list may not describe all possible side effects. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.