

BRENTWOOD PHARMACY
761 Suffolk Ave , Brentwood , NY , 11717
Phone#: 6312733314 Fax#: 6312738863

Drug NDC: 43547-0385-50
Drug Name: GABAPENTIN

How should I use this medicine?

Take this medicine by mouth with a glass of water. Follow the directions on the prescription label. You can take it with or without food. If it upsets your stomach, take it with food. Take your medicine at regular intervals. Do not take it more often than directed. Do not stop taking except on your doctor's advice. If you are directed to break the 600 or 800 mg tablets in half as part of your dose, the extra half tablet should be used for the next dose. If you have not used the extra half tablet within 28 days, it should be thrown away. A special MedGuide will be given to you by the pharmacist with each prescription and refill. Be sure to read this information carefully each time. Talk to your pediatrician regarding the use of this medicine in children. Special care may be needed.

Overdosage: If you think you have taken too much of this medicine contact a poison control center or emergency room at once.
NOTE: This medicine is only for you. Do not share this medicine with others.

Where should I keep my medicine?

Keep out of reach of children. This medicine may cause accidental overdose and death if it taken by other adults, children, or pets. Mix any unused medicine with a substance like cat litter or coffee grounds. Then throw the medicine away in a sealed container like a sealed bag or a coffee can with a lid. Do not use the medicine after the expiration date. Store at room temperature between 15 and 30 degrees C (59 and 86 degrees F).

NOTE: This sheet is a summary. It may not cover all possible information. If you have questions about this medicine, talk to your doctor, pharmacist, or health care provider.

What should I watch for while using this medicine?

Visit your doctor or health care professional for regular checks on your progress. You may want to keep a record at home of how you feel your condition is responding to treatment. You may want to share this information with your doctor or health care professional at each visit. You should contact your doctor or health care professional if your seizures get worse or if you have any new types of seizures. Do not stop taking this medicine or any of your seizure medicines unless instructed by your doctor or health care professional. Stopping your medicine suddenly can increase your seizures or their severity. Wear a medical identification bracelet or chain if you are taking this medicine for seizures, and carry a card that lists all your medications. You may get drowsy, dizzy, or have blurred vision. Do not drive, use machinery, or do anything that needs mental alertness until you know how this medicine affects you. To reduce dizzy or fainting spells, do not sit or stand up quickly, especially if you are an older patient. Alcohol can increase drowsiness and dizziness. Avoid alcoholic drinks. Your mouth may get dry. Chewing sugarless gum or sucking hard candy, and drinking plenty of water will help. The use of this medicine may increase the chance of suicidal thoughts or actions. Pay special attention to how you are responding while on this medicine. Any worsening of mood, or thoughts of suicide or dying should be reported to your health care professional right away. Women who become pregnant while using this medicine may enroll in the North American Antiepileptic Drug Pregnancy Registry by calling 1-888-233-2334. This registry collects information about the safety of antiepileptic drug use during pregnancy.

What is this medicine?

GABAPENTIN (GA ba pen tin) is used to control partial seizures in adults with epilepsy. It is also used to treat certain types of nerve pain.

This medicine may be used for other purposes; ask your health care provider or pharmacist if you have questions.

What should I tell my health care provider before I take this medicine?

They need to know if you have any of these conditions: kidney disease; suicidal thoughts, plans, or attempt; a previous suicide attempt by you or a family member; an unusual or allergic reaction to gabapentin, other medicines, foods, dyes, or preservatives; pregnant or trying to get pregnant; breast-feeding

What if I miss a dose?

If you miss a dose, take it as soon as you can. If it is almost time for your next dose, take only that dose. Do not take double or extra doses.

What side effects may I notice from receiving this medicine?

Side effects that you should report to your doctor or health care professional as soon as possible: allergic reactions like skin rash, itching or hives, swelling of the face, lips, or tongue; worsening of mood, thoughts or actions of suicide or dying; Side effects that usually do not require medical attention (report to your doctor or health care professional if they continue or are bothersome): constipation; difficulty walking or controlling muscle movements; dizziness; nausea; slurred speech; tiredness; tremors; weight gain

This list may not describe all possible side effects. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.