

BRENTWOOD PHARMACY
761 Suffolk Ave , Brentwood , NY , 11717
Phone#: 6312733314 Fax#: 6312738863

Drug NDC: 50458-0579-10
Drug Name: XARELTO

How should I use this medicine?

Take this medicine by mouth with a glass of water. Follow the directions on the prescription label. Take your medicine at regular intervals. Do not take it more often than directed. Do not stop taking except on your doctor's advice. Stopping this medicine may increase your risk of a blood clot. Be sure to refill your prescription before you run out of medicine. If you are taking this medicine after hip or knee replacement surgery, take it with or without food. If you are taking this medicine for atrial fibrillation, take it with your evening meal. If you are taking this medicine to treat blood clots, take it with food at the same time each day. If you are unable to swallow your tablet, you may crush the tablet and mix it in applesauce. Then, immediately eat the applesauce. You should eat more food right after you eat the applesauce containing the crushed tablet. Talk to your pediatrician regarding the use of this medicine in children. Special care may be needed.

Overdosage: If you think you have taken too much of this medicine contact a poison control center or emergency room at once. NOTE: This medicine is only for you. Do not share this medicine with others.

Where should I keep my medicine?

Keep out of the reach of children. Store at room temperature between 15 and 30 degrees C (59 and 86 degrees F). Throw away any unused medicine after the expiration date.

NOTE: This sheet is a summary. It may not cover all possible information. If you have questions about this medicine, talk to your doctor, pharmacist, or health care provider.

What should I watch for while using this medicine?

Visit your doctor or health care professional for regular checks on your progress. Notify your doctor or health care professional and seek emergency treatment if you develop breathing problems; changes in vision; chest pain; severe, sudden headache; pain, swelling, warmth in the leg; trouble speaking; sudden numbness or weakness of the face, arm or leg. These can be signs that your condition has gotten worse. If you are going to have surgery or other procedure, tell your doctor that you are taking this medicine.

What is this medicine?

RIVAROXABAN (ri va ROX a ban) is an anticoagulant (blood thinner). It is used to treat blood clots in the lungs or in the veins. It is also used after knee or hip surgeries to prevent blood clots. It is also used to lower the chance of stroke in people with a medical condition called atrial fibrillation.

This medicine may be used for other purposes; ask your health care provider or pharmacist if you have questions.

What should I tell my health care provider before I take this medicine?

They need to know if you have any of these conditions: bleeding disorders; bleeding in the brain; blood in your stools (black or tarry stools) or if you have blood in your vomit; history of stomach bleeding; kidney disease; liver disease; low blood counts, like low white cell, platelet, or red cell counts; recent or planned spinal or epidural procedure; take medicines that treat or prevent blood clots; an unusual or allergic reaction to rivaroxaban, other medicines, foods, dyes, or preservatives; pregnant or trying to get pregnant; breast-feeding

What if I miss a dose?

If you take your medicine once a day and miss a dose, take the missed dose as soon as you remember. If you take your medicine twice a day and miss a dose, take the missed dose immediately. In this instance, 2 tablets may be taken at the same time. The next day you should take 1 tablet twice a day as directed.

What side effects may I notice from receiving this medicine?

Side effects that you should report to your doctor or health care professional as soon as possible: allergic reactions like skin rash, itching or hives, swelling of the face, lips, or tongue; back pain; redness, blistering, peeling or loosening of the skin, including inside the mouth; signs and symptoms of bleeding such as bloody or black, tarry stools; red or dark-brown urine; spitting up blood or brown material that looks like coffee grounds; red spots on the skin; unusual bruising or bleeding from the eye, gums, or nose; Side effects that usually do not require medical attention (report to your doctor or health care professional if they continue or are bothersome): dizziness; muscle pain

This list may not describe all possible side effects. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.