

BRENTWOOD PHARMACY
761 Suffolk Ave , Brentwood , NY , 11717
Phone#: 6312733314 Fax#: 6312738863

Drug NDC: 50458-0659-60
Drug Name: ULTRAM

How should I use this medicine?

Take this medicine by mouth with a full glass of water. Follow the directions on the prescription label. You can take it with or without food. If it upsets your stomach, take it with food. Do not take your medicine more often than directed. A special MedGuide will be given to you by the pharmacist with each prescription and refill. Be sure to read this information carefully each time. Talk to your pediatrician regarding the use of this medicine in children. Special care may be needed.

Overdosage: If you think you have taken too much of this medicine contact a poison control center or emergency room at once. NOTE: This medicine is only for you. Do not share this medicine with others.

Where should I keep my medicine?

Keep out of the reach of children. This medicine may cause accidental overdose and death if taken by other adults, children, or pets. Mix any unused medicine with a substance like cat litter or coffee grounds. Then throw the medicine away in a sealed container like a sealed bag or a coffee can with a lid. Do not use the medicine after the expiration date. Store at room temperature between 15 and 30 degrees C (59 and 86 degrees F).

NOTE: This sheet is a summary. It may not cover all possible information. If you have questions about this medicine, talk to your doctor, pharmacist, or health care provider.

What should I watch for while using this medicine?

Tell your doctor or health care professional if your pain does not go away, if it gets worse, or if you have new or a different type of pain. You may develop tolerance to the medicine. Tolerance means that you will need a higher dose of the medicine for pain relief. Tolerance is normal and is expected if you take this medicine for a long time. Do not suddenly stop taking your medicine because you may develop a severe reaction. Your body becomes used to the medicine. This does NOT mean you are addicted. Addiction is a behavior related to getting and using a drug for a non-medical reason. If you have pain, you have a medical reason to take pain medicine. Your doctor will tell you how much medicine to take. If your doctor wants you to stop the medicine, the dose will be slowly lowered over time to avoid any side effects. There are different types of narcotic medicines (opiates). If you take more than one type at the same time or if you are taking another medicine that also causes drowsiness, you may have more side effects. Give your health care provider a list of all medicines you use. Your doctor will tell you how much medicine to take. Do not take more medicine than directed. Call emergency for help if you have problems breathing or unusual sleepiness. You may get drowsy or dizzy. Do not drive, use machinery, or do anything that needs mental alertness until you know how this medicine affects you. Do not stand or sit up quickly, especially if you are an older patient. This reduces the risk of dizzy or fainting spells. Alcohol can increase or decrease the effects of this medicine. Avoid alcoholic drinks. You may have constipation. Try to have a bowel movement at least every 2 to 3 days. If you do not have a bowel movement for 3 days, call your doctor or health care professional. Your mouth may get dry. Chewing sugarless gum or sucking hard candy, and drinking plenty of water may help. Contact your doctor if the problem does not go away or is severe.

What is this medicine?

TRAMADOL (TRA ma dole) is a pain reliever. It is used to treat moderate to severe pain in adults.

This medicine may be used for other purposes; ask your health care provider or pharmacist if you have questions.

What should I tell my health care provider before I take this medicine?

They need to know if you have any of these conditions: brain tumor, depression, drug abuse or addiction, head injury, if you frequently drink alcohol containing drinks, kidney disease or trouble passing urine, liver disease, lung disease, asthma, or breathing problems, seizures or epilepsy, suicidal thoughts, plans, or attempt; a previous suicide attempt by you or a family member, an unusual or allergic reaction to tramadol, codeine, other medicines, foods, dyes, or preservatives, pregnant or trying to get pregnant, breast-feeding

What if I miss a dose?

If you miss a dose, take it as soon as you can. If it is almost time for your next dose, take only that dose. Do not take double or extra doses.

What side effects may I notice from receiving this medicine?

Side effects that you should report to your doctor or health care professional as soon as possible: allergic reactions like skin rash, itching or hives, swelling of the face, lips, or tongue; breathing problems; confusion; seizures; signs and symptoms of low blood pressure like dizziness; feeling faint or lightheaded, falls; unusually weak or tired; trouble passing urine or change in the amount of urine; Side effects that usually do not require medical attention (report to your doctor or health care professional if they continue or are bothersome): constipation; dry mouth; nausea, vomiting; tiredness

This list may not describe all possible side effects. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.