

BRENTWOOD PHARMACY
761 Suffolk Ave , Brentwood , NY , 11717
Phone#: 6312733314 Fax#: 6312738863

Drug NDC: 58914-0018-10
Drug Name: ULTRASE MT18

How should I use this medicine?

Take this medicine by mouth with a glass of water. Follow the directions on the prescription label. Take with food. Do not crush or chew the contents of the capsule. If you or your child have trouble swallowing, you may open the capsule and sprinkle the contents on soft foods that do not require chewing such as applesauce, pureed bananas, or pears. Swallow the mixture right away followed with water or juice. Do not store the mixture. If you are giving this medicine to an infant, you may sprinkle the contents directly into your child's mouth. Give the medicine right before each feeding of formula or breast milk. Do not mix capsule contents directly into formula or breast milk. Make sure the medicine is swallowed completely and that no medicine is left in the mouth. Take your doses at regular intervals. Do not take your medicine more often than directed. A special MedGuide will be given to you by the pharmacist with each prescription and refill of delayed-release capsules (Creon, Zenpep, or Pancreaze). Be sure to read this information carefully each time. Talk to your pediatrician regarding the use of this medicine in children. While this medicine may be prescribed for children for selected conditions precautions do apply.

Overdosage: If you think you have taken too much of this medicine contact a poison control center or emergency room at once. NOTE: This medicine is only for you. Do not share this medicine with others.

Where should I keep my medicine?

Keep out of the reach of children. Store at room temperature between 15 and 25 degrees C (59 and 77 degrees F). Do not refrigerate. Protect from moisture. Throw away any unused medicine after the expiration date.

NOTE: This sheet is a summary. It may not cover all possible information. If you have questions about this medicine, talk to your doctor, pharmacist, or health care provider.

What should I watch for while using this medicine?

Visit your doctor for regular check ups. Talk to your doctor before you change brands of this medicine. Each brand has different amounts of enzymes. You may need to be on a special diet while taking this medicine. Also, ask your doctor how much water you need to drink. This medicine may increase your chance of having a rare bowel disorder. The risk of having this condition may be reduced by following the dosing directions that your healthcare professional gives you. Call your healthcare professional right away if you have any unusual or severe stomach pain. This medicine may increase blood uric acid levels, for example, worsening of gout, or painful, swollen joints. Call your healthcare professional right away if you have any of these symptoms. Be careful if you open the capsule. This medicine can irritate the lungs if you breathe it in. Also, do not hold the medicine in your mouth or chew it. This may cause mouth sores. This medicine may affect blood sugar levels. If you have diabetes, check with your doctor or health care professional before you change your diet or the dose of your diabetic medicine. Women should inform their doctor if they wish to become pregnant or think they might be pregnant.

What is this medicine?

PANCRELIPASE (pan cre LI pase) helps to improve digestion of food by replacing digestive enzymes. This medicine is used to treat health conditions that cause your body to produce less of these enzymes.

This medicine may be used for other purposes; ask your health care provider or pharmacist if you have questions.

What should I tell my health care provider before I take this medicine?

They need to know if you have any of these conditions: a history of intestinal blockage or a condition called 'fibrosing colonopathy' abnormally high uric acid in the blood diabetes gout kidney disease trouble swallowing capsules an unusual or allergic reaction to pancrelipase, pancreatin, pork, pork protein, other medicines, foods, dyes, or preservatives pregnant or trying to get pregnant breast-feeding

What if I miss a dose?

If you miss a dose, take it as soon as you can. If it is almost time for your next dose, take only that dose. Do not take double or extra doses.

What side effects may I notice from receiving this medicine?

Side effects that you should report to your doctor or health care professional as soon as possible: allergic reactions like skin rash, itching or hives, swelling of the face, lips, or tongue fever or chills, sore throat severe stomach pain or bloating shortness of breath skin rash trouble passing stool vomiting Side effects that usually do not require medical attention (report to your doctor or health care professional if they continue or are bothersome): constipation or diarrhea cough dizziness headache nausea stomach gas stomach pain weight loss

This list may not describe all possible side effects. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.