

BRENTWOOD PHARMACY
761 Suffolk Ave , Brentwood , NY , 11717
Phone#: 6312733314 Fax#: 6312738863

Drug NDC: 59743-0002-01
Drug Name: G-PHED

How should I use this medicine?

Take this medicine by mouth with a full glass of water. Follow the directions on the prescription label. Do not crush or chew. Take your doses at regular times. Do not take more medicine than directed. Talk to your pediatrician regarding the use of this medicine in children. While this drug may be prescribed for children as young as 2 years old for selected conditions, precautions do apply. Patients over 60 years old may have a stronger reaction to this medicine and need smaller doses.

Overdosage: If you think you have taken too much of this medicine contact a poison control center or emergency room at once. NOTE: This medicine is only for you. Do not share this medicine with others.

Where should I keep my medicine?

Keep out of the reach of children. Store at room temperature between 15 and 30 degrees C (59 and 86 degrees F). Protect from heat and moisture. Throw away any unused medicine after the expiration date.

NOTE: This sheet is a summary. It may not cover all possible information. If you have questions about this medicine, talk to your doctor, pharmacist, or health care provider.

What should I watch for while using this medicine?

Tell your doctor or health care professional if your symptoms do not improve within 5 days or if they get worse. If you have trouble falling asleep at night, take the last dose of the day at least a few hours before bedtime. Drink water as instructed while you are taking this medicine to help loosen mucus. Your mouth may get dry. Chewing sugarless gum or sucking hard candy, and drinking plenty of water may help. Contact your doctor if the problem does not go away or is severe. This medicine may cause dry eyes and blurred vision. If you wear contact lenses you may feel some discomfort. Lubricating drops may help. See your eye doctor if the problem does not go away or is severe.

What is this medicine?

GUAIFENESIN; PSEUDOEPHEDRINE (gwye FEN e sin; soo doe e FED rin) is a decongestant, expectorant combination. It is used to provide relief from cough and congestion. This medicine will not treat an infection.

This medicine may be used for other purposes; ask your health care provider or pharmacist if you have questions.

What should I tell my health care provider before I take this medicine?

They need to know if you have any of the following conditions: blood vessel disease, diabetes, fever, glaucoma, heart disease or heart rhythm problems, high blood pressure, other chronic disease, over active thyroid, prostate trouble, an unusual or allergic reaction to guaifenesin, pseudoephedrine, other medicines, foods, dyes, or preservatives, pregnant or trying to get pregnant, breast-feeding

What if I miss a dose?

If you miss a dose, take it as soon as you can. If it is almost time for your next dose, take only that dose. Do not take double or extra doses.

What side effects may I notice from receiving this medicine?

Side effects that you should report to your doctor or health care professional as soon as possible: allergic reactions like skin rash, itching or hives, swelling of the face, lips, or tongue; confusion; difficulty passing urine; fast or irregular heartbeat; hallucinations; high blood pressure; seizure; tremor; trouble breathing; Side effects that usually do not require medical attention (report to your doctor or health care professional if they continue or are bothersome): anxiety, excitement, irritability, nervousness; diarrhea; dizziness; headache; loss of appetite; stomach upset, nausea

This list may not describe all possible side effects. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.