

**BRENTWOOD PHARMACY**  
**761 Suffolk Ave , Brentwood , NY , 11717**  
**Phone#: 6312733314 Fax#: 6312738863**

**Drug NDC:** 61451-5016-07  
**Drug Name:** B & O16-A SUPPRETTE

#### **How should I use this medicine?**

This medicine is for rectal use only. Do not take by mouth. Wash your hands before and after use. Take off the foil wrapping. Wet the tip of the suppository with cold tap water to make it easier to use. Lie on your side with your lower leg straightened out and your upper leg bent forward toward your stomach. Lift upper buttock to expose the rectal area. Apply gentle pressure to insert the suppository completely into the rectum, pointed end first. Hold buttocks together for a few seconds. Remain lying down for about 15 minutes to avoid having the suppository come out. Do not use more often than directed. A special MedGuide will be given to you by the pharmacist with each prescription and refill. Be sure to read this information carefully each time. Talk to your pediatrician regarding the use of this medicine in children. While this drug may be prescribed for children as young as 13 years old for selected conditions, precautions do apply.

**Overdosage: If you think you have taken too much of this medicine contact a poison control center or emergency room at once.**  
**NOTE: This medicine is only for you. Do not share this medicine with others.**

#### **Where should I keep my medicine?**

Keep out of the reach of children. This medicine can be abused. Keep your medicine in a safe place to protect it from theft. Do not share this medicine with anyone. Selling or giving away this medicine is dangerous and against the law. Store at room temperature between 15 and 30 degrees C (59 and 86 degrees F). Do NOT refrigerate. Protect from moisture. Throw away any unused medicine after the expiration date.

**NOTE: This sheet is a summary. It may not cover all possible information. If you have questions about this medicine, talk to your doctor, pharmacist, or health care provider.**

#### **What should I watch for while using this medicine?**

Tell your doctor or health care professional if your pain does not go away, if it gets worse, or if you have new or a different type of pain. You may develop tolerance to the medicine. Tolerance means that you will need a higher dose of the medicine for pain relief. Tolerance is normal and is expected if you take the medicine for a long time. Do not suddenly stop taking your medicine because you may develop a severe reaction. Your body becomes used to the medicine. This does NOT mean you are addicted. Addiction is a behavior related to getting and using a drug for a non-medical reason. If you have pain, you have a medical reason to take pain medicine. Your doctor will tell you how much medicine to take. If your doctor wants you to stop the medicine, the dose will be slowly lowered over time to avoid any side effects. There are different types of narcotic medicines (opiates). If you take more than one type at the same time or if you are taking another medicine that also causes drowsiness, you may have more side effects. Give your health care provider a list of all medicines you use. Your doctor will tell you how much medicine to take. Do not take more medicine than directed. Call emergency for help if you have problems breathing or unusual sleepiness. You may get drowsy or dizzy. Do not drive, use machinery, or do anything that needs mental alertness until you know how this medicine affects you. Do not stand or sit up quickly, especially if you are an older patient. This reduces the risk of dizzy or fainting spells. Alcohol may interfere with the effect of this medicine. Avoid alcoholic drinks. This medicine may cause constipation. Try to have a bowel movement at least every 2 to 3 days. If you do not have a bowel movement for 3 days, call your doctor or health care professional. Your mouth may get dry. Chewing sugarless gum or sucking hard candy, and drinking plenty of water may help. Contact your doctor if the problem does not go away or is severe.

#### **What is this medicine?**

BELLADONNA; OPIUM (bell a DON a; OH-pee-uhm) is a combination of two different drugs. It treats pain associated from spasm that may occur after surgery of the urinary tract.

**This medicine may be used for other purposes; ask your health care provider or pharmacist if you have questions.**

#### **What should I tell my health care provider before I take this medicine?**

They need to know if you have any of these conditions: drug abuse or addiction; glaucoma; head injury; heart disease; high blood pressure; if you frequently drink alcohol-containing drinks; intestinal disease; kidney disease; liver disease; myasthenia gravis; lung or breathing disease, like asthma; prostate trouble; seizures; an unusual or allergic reaction to belladonna, opium, other medicines, foods, dyes, or preservatives; pregnant or trying to get pregnant; breast-feeding

#### **What if I miss a dose?**

If you miss a dose, take it as soon as you can. If it is almost time for your next dose, take only that dose. Do not take double or extra doses.

#### **What side effects may I notice from receiving this medicine?**

Side effects that you should report to your doctor or health care professional as soon as possible: allergic reactions like skin rash, itching or hives, swelling of the face, lips, or tongue; breathing problems; confusion; signs and symptoms of low blood pressure like dizziness; feeling faint or lightheaded, falls; unusually weak or tired; trouble passing urine or change in the amount of urine; Side effects that usually do not require medical attention (report to your doctor or health care professional if they continue or are bothersome): constipation; dry mouth; nausea, vomiting; tiredness

**This list may not describe all possible side effects. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.**