

**BRENTWOOD PHARMACY**  
**761 Suffolk Ave , Brentwood , NY , 11717**  
**Phone#: 6312733314 Fax#: 6312738863**

**Drug NDC: 61958-1004-01**  
**Drug Name: RANEXA**

**How should I use this medicine?**

Take this medicine by mouth with a glass of water. Follow the directions on the prescription label. Do not cut, crush, or chew this medicine. Take with or without food. Do not take this medication with grapefruit juice. Take your doses at regular intervals. Do not take your medicine more often than directed.  
Talk to your pediatrician regarding the use of this medicine in children. Special care may be needed.

**Overdosage: If you think you have taken too much of this medicine contact a poison control center or emergency room at once.**  
**NOTE: This medicine is only for you. Do not share this medicine with others.**

**Where should I keep my medicine?**

Keep out of the reach of children.  
Store at room temperature between 15 and 30 degrees C (59 and 86 degrees F). Throw away any unused medicine after the expiration date.

**NOTE: This sheet is a summary. It may not cover all possible information. If you have questions about this medicine, talk to your doctor, pharmacist, or health care provider.**

**What should I watch for while using this medicine?**

Visit your doctor for regular check ups. Tell your doctor or healthcare professional if your symptoms do not start to get better or if they get worse. This medicine will not relieve an acute attack of angina or chest pain.  
This medicine can change your heart rhythm. Your health care provider may check your heart rhythm by ordering an electrocardiogram (ECG) while you are taking this medicine.  
You may get drowsy or dizzy. Do not drive, use machinery, or do anything that needs mental alertness until you know how this medicine affects you. Do not stand or sit up quickly, especially if you are an older patient. This reduces the risk of dizzy or fainting spells. Alcohol may interfere with the effect of this medicine. Avoid alcoholic drinks.  
If you are scheduled for any medical or dental procedure, tell your healthcare provider that you are taking this medicine. This medicine can interact with other medicines used during surgery.

**What is this medicine?**

RANOLAZINE (ra NOE la zeen) is a heart medicine. It is used to treat chronic chest pain (angina). This medicine must be taken regularly. It will not relieve an acute episode of chest pain.

**This medicine may be used for other purposes; ask your health care provider or pharmacist if you have questions.**

**What should I tell my health care provider before I take this medicine?**

They need to know if you have any of these conditions:  
heart disease  
irregular heartbeat  
kidney disease  
liver disease  
low levels of potassium or magnesium in the blood  
an unusual or allergic reaction to ranolazine, other medicines, foods, dyes, or preservatives  
pregnant or trying to get pregnant  
breast-feeding

**What if I miss a dose?**

If you miss a dose, take it as soon as you can. If it is almost time for your next dose, take only that dose. Do not take double or extra doses.

**What side effects may I notice from receiving this medicine?**

Side effects that you should report to your doctor or health care professional as soon as possible:  
allergic reactions like skin rash, itching or hives, swelling of the face, lips, or tongue  
breathing problems  
changes in vision  
fast, irregular or pounding heartbeat  
feeling faint or lightheaded, falls  
low or high blood pressure  
numbness or tingling feelings  
ringing in the ears  
tremor or shakiness  
slow heartbeat (fewer than 50 beats per minute)  
swelling of the legs or feet  
Side effects that usually do not require medical attention (report to your doctor or health care professional if they continue or are bothersome):  
constipation  
drowsy  
dry mouth  
headache  
nausea or vomiting  
stomach upset

**This list may not describe all possible side effects. Call your doctor for medical advice about side effects.**  
**You may report side effects to FDA at 1-800-FDA-1088.**